

PLEASE USE THE FOLLOWING INFORMATION WHEN COMPLETING YOUR DONOR HISTORY FORM.

SOME MEDICATIONS MAY AFFECT YOUR ELIBILITY TO DONATE BLOOD. PLEASE TELL US IF YOU...

ARE BEING TREATED WITH ANY OF THE FOLLOWING TYPES OF MEDICATIONS:	OR HAVE TAKEN:	WHICH IS ALSO CALLED:	ANYTIME IN THE LAST:	
Anti-platelet agents: Usually taken to prevent stroke or heart attack (deferral applies only to Platelet donations via apheresis)	Feldene	piroxicam	2 Days	
	Effient	prasugrel	3 Days	
	Brilinta	ticagrelor	7 Days	
	Plavix	clopidogrel	14 Days	
	Ticlid	ticlopidine		
	Zontivity	vorapaxar	1 Month	
Anticoagulants or "blood thinners": Usually taken to prevent blood clots in the legs and lungs and to prevent strokes (deferral applies only to Platelet and/or Plasma donations via apheresis)	Arixtra	fondaparinux	2 Days	
	Eliquis	apixaban		
	Fragmin	dalteparin		
	Lovenox	enoxaparin		
	Pradaxa	dabigatran		
	Savaysa	edoxaban		
	Xarelto	rivaroxaban		
	Coumadin, Warfilone, Jantoven	warfarin	7 Days	
Heparin, low molecular weight heparin				
Acne treatment	Accutane Amnesteem Absorica Claravis Myorisan Sotret Zenatane	Isotretinoin	1 Month	
Multiple myeloma	Thalomid Revlimid	thalidomide lenalidomide		
Rheumatoid arthritis	Rinvoq	upadacitinib		
Hair loss remedy	Propecia	finasteride		
Prostate symptoms	Proscar	finasteride		
	Avodart Jalyn	dutasteride	6 Months	
Immunosuppressant	Cellcept	mycophenolate mofetil	6 Weeks	
Hepatitis exposure	Hepatitis B Immune Globulin	HBIG	3 Months	
HIV prevention (Also known as PrEP or PEP)	Any medication taken by mouth (oral) to prevent HIV	Truvada		emtricitabine and tenofovir disoproxil fumarate
		Descovy		emtricitabine and tenofovir alafenamide
	Injectable HIV prevention	Apretude	cabotegravir	2 Years
Basal cell skin cancer	Erivedge Odomzo	vismodegib sonidegib	2 Years	
Relapsing multiple sclerosis	Aubagio	teriflunomide		
Rheumatoid arthritis	Arava	leflunomide		
Psoriasis	Soriatane	acitretin	3 years	
	Tegison	etretinate	Ever	
HIV treatment	Any medication to treat HIV. May also be called antiretroviral therapy (ART)			
Experimental medication			As defined by the medical director	

Donors SHOULD NOT discontinue medications prescribed or recommended by their physician in order to donate blood.